

Avoiding barren plateaus via transferability of smooth solutions in Hamiltonian Variational Ansatz

Antonio A. Mele, Glen B. Mbeng, Giuseppe E. Santoro, Mario Collura, Pietro Torta


arXiv:2206.01982



QuAntonioMele



antonioannamele.com



Antonio Anna Mele, Freie Universität Berlin

OUTLINE

- **Variational Quantum Algorithms and Barren Plateaus**
- **QAOA-inspired ansatz**
- **Pattern of optimal parameters**
- **Solution transferability**
- **Avoiding Barren Plateaus**

Variational Quantum Algorithms

- Leading **NISQ** strategy
- The problem is encoded in **minimising a cost function**

(e.g. finding Ground state of an Hamiltonian)


(e.g. Hamiltonian expectation value)

- The **main steps** are:

1. State preparation using a **parameterized circuit**
2. **Measurement** process
3. Classical **optimization**

MAIN DIFFICULTIES:


- **Noise**
- **Non-convex** optimization
- Flat landscape (a.k.a. **Barren Plateaus**)



VQAs scheme [Bittel et al., PRL (2021)]

Barren Plateaus


High circuit expressibility




Exponential vanishing gradients
with number of qubits N
(Barren Plateaus definition)

[McClean et al., Nat. Comm. (2018)]
[Holmes et al., PRX Q. (2022)]


Estimation accuracy at least exponential



Exp. number of measurements needed



💀 Serious problem for VQAs 💀



Orange (blue) landscape $N = 24$
 $(N = 4)$ qubits. [Cerezo et al., Nature(2021)]

Problem-inspired Ansatz

$$|\psi(\gamma)\rangle = \prod_{m=1}^P e^{-i\gamma_{m,M}H_M} \dots e^{-i\gamma_{m,1}H_1} |\psi_0\rangle$$

The problem **Hamiltonian** is a **linear combination of these generators**

Easy-to-prepare state

These are **symmetry-preserving** ansatz, known as **Hamiltonian Variational Ansatz** (generalisation of QAOA)

We analyzed:

$$H_{XYZ} = \sum_{j=1}^N \left(X_j X_{j+1} + \Delta_Y Y_j Y_{j+1} + \Delta_Z Z_j Z_{j+1} \right)$$


$$H_{\text{LTFIM}} = \sum_{j=1}^N Z_j Z_{j+1} - g_x \sum_{j=1}^N X_j - g_z \sum_{j=1}^N Z_j$$

Although **symmetry-ansatz**,
there can be Barren Plateaus

[Larocca et al., ArXiv (2021)]

Pattern in Optimal Parameters


$$(\alpha_1, \dots, \alpha_P, \beta_1, \dots, \beta_P)$$





How can we find this pattern?

INTERP strategy


[Zhou et al., PRX. (2018)]




Warm-start for the optimization when you go from $P \rightarrow P+1$



Transferability of solutions



The warm-start allows to avoid the flat region



OPEN QUESTIONS

- 2D systems
- This helps avoiding bad local minima and BPs, but what about noise resilience?
- Analytical understanding (connection with Adiabatic QC?)

**THANKS FOR YOUR
ATTENTION!**